

ZIKA VIRUS UPDATE **2016**

LARRY M. BUSH, MD, FACP

Affiliated Professor of Clinical Biomedical Sciences

Charles E. Schmidt College of Medicine

Florida Atlantic University

Affiliate Associate Professor of Medicine

University of Miami – Miller School of Medicine / Palm Beach County



Early Optimism about Antimicrobials



William H. Stewart, U.S. Surgeon General 1965-1969, is purported to have said, “We have closed the book on infectious disease.”

The End of Infectious Disease

- Since 1967:
 - Legionnaire's disease
 - Toxic shock syndrome
 - AIDS
 - Lyme disease
 - West Nile encephalitis
 - SARS
 - Avian Flu
- Chronic diseases associated with pathogens:
 - Peptic ulcers (*Helicobacter pylori*)
 - Liver cancer (Hepatitis B and C)
 - Lyme arthritis (*Borrelia burgdorferi*)

NEWLY IDENTIFIED INFECTIOUS DISEASES AND PATHOGENS (2)

<u>Year</u>	<u>Disease or Pathogen</u>
2012	MERS-CoV
2009	H1N1
2004	Avian influenza (human cases)
2003	SARS
1999	Nipah virus
1997	H5N1 (avian influenza A virus)
1996	New variant Creutzfeldt-Jacob disease; Australian bat lyssavirus
1995	Human herpesvirus 8 (Kaposi's sarcoma virus)
1994	Sabia virus; Hendra virus


NEWLY IDENTIFIED INFECTIOUS DISEASES AND PATHOGENS (2)

<u>Year</u>	<u>Disease or Pathogen</u>
2012	MERS-CoV
2009	H1N1
2004	Avian influenza (human cases)
2003	SARS
1999	Nipah virus
1997	H5N1 (avian influenza A virus)
1996	New variant Creutzfeldt-Jacob disease; Australian bat lyssavirus
1995	Human herpesvirus 8 (Kaposi's sarcoma virus)
1994	Sabia virus; Hendra virus

Emerging and Re-emerging Infectious Diseases

- **Emerging infectious diseases:** Infectious diseases that have newly appeared in a population.
- Global :
- Regional:
- **Re-emerging Diseases:** Diseases' incidence in human has increased during the last 20 years or threatens to increase in the near future.
- Global:
- Regional:

GLOBAL EXAMPLES OF EMERGING AND RE-EMERGING INFECTIOUS DISEASES



Factors responsible for emerging of infections.

I-Ecological changes and Agricultural development.



Placing the people in contact with a natural reservoir or host of a hitherto unfamiliar, but usually already present,

Example 1 :

Reforestation in USA

```
graph TD; A[Reforestation in USA] --> B[Increased the number of deer & deer ticks]; B --> C[Increased Human contact with deers]; B --> D[Deer ticks are natural reservoir of Lyme diseases]; C --> E[Human affliction by Lyme disease]; D --> E;
```

Increased the number of deer & deer ticks

Increased Human contact with deers

Deer ticks are natural reservoir of Lyme diseases

Human affliction by Lyme disease

II- Changes in Human demographics and behaviours

Inflation of population size



Insufficient infrastructures



Use open containers for water




Breeding mosquitos



Dengue fever in Asia

Growth in More, Less Developed Countries

Billions



Source: United Nations, *World Population Prospects: The 2004 Revision* (medium scenario), 2005.

- **Urbanization-more people concentrated in cities-often without adequate infrastructure**
- **Increases in the elderly populations**
- **Increases in children in daycare: working woman with kids under 5 was 30% in 1970, is 75% in 2000.**
- **Fast paced Lifestyles- increase in convenience items and more stress**
- **High-risk behavior- Drug use and unprotected sex**

War and Famine

War refugees are a full 1% of the global population

War refugees are forced onto new areas where they are exposed to new microbes from vectors and people.

War and famine are closely linked.

In 2001, tracking 16 countries with “food emergencies”, showed that 9 were because of civil unrest.

Famine is also caused by social, economic, and political forces. Weather and HIV/AIDS

- **III-International travel and Commerce.**
- **365 days to circumnavigate the globe...now it takes 36 hours**
- **-used to quarantine ships, but 36 h faster than disease incubation**
- **400 million people per year travel internationally**
- **increased incidence of both Tuberculosis and Influenza transmission on long flights**


The global air network



Commercial Air Traffic Over a 24 Hour Period



Average annual number of global airline passengers by decade, 1950-2010



- -Transportation of products is an increased concern.
- -rapid transport of disease harboring fresh products.
- -transport of livestock facilitates movements of viruses and arthropods (especially ticks)

Travel and HIV/AIDS.

- Silk route and plague.
- Slaves trade and yellow fever.
- Migration to new world and smallpox.

Cholera and Hajj.

IV- Technology and industry

- Modern mass production increased the chance of accidental contamination and amplifies the effect of such contamination.
 - Contamination of hamburger meat by E.coli strains causing haemolytic uraemic syndrome.
 - Feeding cattle by byproducts of sheep causing bovine spongiform encephalitis.

VI- Breakdown of public health measures.

- -Decrease in chlorine in water supplies lead to rapid spread of cholera in South America.
- Non functioning water plant in Wisconsin, USA lead to outbreak of waterborne cryptosporidium.

Major Factors Contributing to the Emergence of Infectious Diseases

- 1. Human demographics and behavior**
- 2. Technology and industry**
- 3. Economic development and land use**
- 4. International travel and commerce**
- 5. Microbial adaptation and change**
- 6. Breakdown of public health measures**

Institute of Medicine Report 1992



THE EMERGENCE OF ZIKA VIRUS

1947 – Isolation from Rhesus monkey, Zika forest, Uganda

1948 – Isolated from *Aedes africanus* mosquitos, Zika forest, 6% people in area test positive serologically

1952 – Nigeria, isolated from child, fever & headache

1956 – Human volunteer inoculation, limited febrile illness

1962 – Uganda, person with rash, fever, body pain

1960's – 1970's _ Africa and Asia febrile illness, sero +, Nigeria sero-prevalence 48% to 56%

RECENT ZIKA VIRUS EPIDEMICS

**2007 – Yap Island, Micronesia (pop. 10,000),
rash, fever, arthralgia, conjunctivitis, 73%
inhabitants IgM antibodies, 19% clinical illness**

**2013 – French Polynesia islands (pop. 270,000),
28,000 cases, Guillain-Barre Syndrome (20% up)**

**2015 – NE Brazil, Americas & Caribbean, 33
countries**

**Feb 2016 – WHO microcephaly & neuro
disorders (GBS), 4th international emergency**

Rates of Microcephaly by State in Brazil

2010-2014 vs 2015-2016

Countries and territories with confirmed cases of Zika virus (autochthonous transmission), 2015-2016 & Rates of microcephaly by state in Brazil, 2010-2014 and 2015-2016




Pan American Health Organization



World Health Organization
Regional Office for the Americas

Updated as of Epidemiological Week 2 (January 10-16, 2016)



Microcephaly rates by state in Brazil (cases per 1,000 live births)

Rate

- 0.1 - 1.0
- 1.1 - 15.0
- 15.1 - 30.0
- 30.1 - 45.0
- 45.1 - 100.1

□ Countries

Countries with Zika confirmed cases

- EW 2, 2016
- Country limits
- Brazil state boundaries



Data Source:
Reported from the IHR National Focal Points and through the Ministry of Health websites.

Map Production:
PAHO-WHO AD CHA IR AR O



Zika epidemic and GBS

Countries reporting increased incidence of Guillain-Barre Syndrome and Zika prevalence:

- **Brazil**
- **El Salvador**
- **Columbia**
- **Suriname**
- **Venezuela**
- Media reports in Puerto Rico and Honduras have not been confirmed
- Martinique has reported no increase of GBS,

Zika virus (Zika)

- Single stranded RNA virus
- Genus *Flavivirus*, family *Flaviviridae*
- Closely related to dengue, yellow fever, Japanese encephalitis, and West Nile viruses
- Primarily transmitted through the bite of an infected *Aedes* species mosquito (*Ae. aegypti* and *Ae. albopictus*).



Aedes aegypti



Aedes albopictus

The Vector



Aedes Aegypti

- Daytime mosquito – dawn and dusk; however due to LED light use is thought to be active at night
- Limited movement – flying radius of 500m
- Female lays ~80 eggs every 3 days
- Needs blood meal to lay eggs
- Eggs can survive several years out of water
- 4 stages of the lifecycle, 3 are water related
- Well adapted to human habitats (water storage, waste, white lights, air conditioning, bromeliads)

ZIKA VIRUS EPIDEMIOLOGY


A solid horizontal bar in a light cyan color, spanning the width of the slide, positioned below the title.

Where has Zika virus been found?

- Before 2015, Zika outbreaks occurred in Africa, Southeast Asia, and the Pacific Islands.
- Currently outbreaks are occurring in many countries and territories.



Possible Geographic Distribution of *Aedes aegypti*



ALL COUNTRIES & TERRITORIES with ACTIVE ZIKA VIRUS TRANSMISSION (as of August 2, 2016)



Case Counts in the US

As of [August 3, 2016](#) (5 am EST)

US States

- Locally acquired mosquito-borne cases reported: 6
- Travel-associated cases reported: 1,818
- Laboratory acquired cases reported: 1
- Total: 1,825
 - Sexually transmitted: 16
 - Guillain-Barré syndrome: 5

MAPS OF ZIKA IN THE US

[More](#)

US Territories

- Locally acquired cases reported: 5,525
- Travel-associated cases reported: 23
- Total: 5,548*
 - Guillain-Barré syndrome: 18
 -

*Sexually transmitted cases are not reported for areas with local mosquito-borne transmission of Zika virus because it is not possible to determine whether infection occurred due to mosquito-borne or sexual transmission.

Department of Health

Daily Zika Update

By Florida Department of Health, Office
of Communications

August 05, 2016

Infection Type	Infection Count
Travel-Related Infections of Zika	351
Non-Travel Related Infections of Zika	16
Infections Involving Pregnant Women	55

Laboratory-confirmed Zika virus disease cases reported to ArboNET by state or territory — United States, 2015–2016 (as of August 3, 2016)

State	Travel-associated cases* No. (% of cases in states) (N=1,819)	Locally acquired cases† No. (% of cases in states) (N=6)
Alabama	11 (1)	0 (0)
Arizona	10 (1)	0 (0)
Arkansas	5 (<1)	0 (0)
California	87 (5)	0 (0)
Colorado	19 (1)	0 (0)
Connecticut	39 (2)	0 (0)
Delaware	10 (1)	0 (0)
District of Columbia	10 (1)	0 (0)
Florida	322 (18)	6 (100)
Georgia	42 (2)	0 (0)
Hawaii	10 (1)	0 (0)
Illinois	33 (2)	0 (0)
Indiana	20 (1)	0 (0)
Iowa	9 (1)	0 (0)
Kansas	8 (<1)	0 (0)
Kentucky		0 (0)
Louisiana	9 (<1)	0 (0)
Maine	9 (<1)	0 (0)
Maryland	54 (3)	0 (0)
Massachusetts	54 (3)	0 (0)
Michigan	17 (1)	0 (0)
Minnesota	29 (2)	0 (0)
Mississippi	14 (1)	0 (0)
Missouri	10 (<1)	0 (0)
Montana	1 (<1)	0 (0)
Nebraska	4 (<1)	0 (0)
Nevada	11 (1)	0 (0)
New Hampshire	8 (<1)	0 (0)
New Jersey	50 (3)	0 (0)
New Mexico	3 (<1)	0 (0)
New York	491 (27)	0 (0)
North Carolina	30 (1)	0 (0)
North Dakota	1 (<1)	0 (0)
Ohio	26 (1)	0 (0)
Oklahoma	13 (1)	0 (0)
Oregon	14 (1)	0 (0)
Pennsylvania††	58 (3)	0 (0)
Rhode Island	18 (1)	0 (0)
South Carolina	28 (2)	0 (0)
Tennessee	22 (1)	0 (0)
Texas	89 (5)	0 (0)
Utah	6 (<1)	0 (0)
Vermont	6 (<1)	0 (0)
Virginia	57 (3)	0 (0)
Washington	16 (1)	0 (0)
West Virginia	9 (1)	0 (0)
Wisconsin	17 (1)	0 (0)
Territories	Travel-associated cases* No. (% of cases in territories) (N=23)	Locally acquired cases† No. (% of cases in territories) (N=5,525)
American Samoa	0 (0)	44 (1)
Puerto Rico	22 (96)	5,460 (99)
US Virgin Islands	1 (4)	21 (<1)

*Travelers returning from affected areas, their sexual contacts, or infants infected in utero

†Presumed local mosquito-borne transmission

†† One additional case acquired through laboratory transmission

Pregnant Women with Any Laboratory Evidence of Possible Zika Virus Infection

US States and the District of Columbia

479

***Includes aggregated data reported to the US Zika Pregnancy Registry as of July 28, 2016**

US Territories**

493

****Includes aggregated data from the US territories reported to the US Zika Pregnancy Registry and data from Puerto Rico reported to the Zika Active Pregnancy Surveillance System as of July 28, 2016**

Pregnancy Outcomes in the United States and the District of Columbia

Liveborn infants with birth defects*

15

Includes aggregated data reported to the [US Zika Pregnancy Registry as of July 28, 2016](#)

Pregnancy losses with birth defects**

6

Includes aggregated data reported to the [US Zika Pregnancy Registry as of July 28, 2016](#)

Pregnancy Outcomes in the United States Territories

Liveborn infants with birth defects*

0

Includes aggregated data from the US territories reported to the [US Zika Pregnancy Registry and data from Puerto Rico reported to the Zika Active Pregnancy Surveillance System as of July 28, 2016](#)

Pregnancy losses with birth defects**

1

Includes aggregated data from the US territories reported to the [US Zika Pregnancy Registry and data from Puerto Rico reported to the Zika Active Pregnancy Surveillance System as of July 28, 2016](#)

🌸 Zika outbreak prompts travel warning for area of Miami

🌸 3:32 a.m. Tuesday, Aug. 2, 2016 | Filed in: News

ARE YOU SURPRISED ?

It's here: Florida 1st state with local Zika cases, transmitted by mosquito



Palm Beach County has first case of locally- acquired Zika virus

7:16 p.m. Monday, Aug. 8, 2016 | Filed in: News

Modes of transmission

- Bite from an infected mosquito
- Maternal-fetal
 - Intrauterine
 - Perinatal
- Sexual transmission from infected male partners
- Laboratory exposure
- Theoretical: blood transfusion, organ and tissue transplant, fertility treatment, and breast feeding



Example Zika virus incidence and attack rates, Yap 2007

- Infection rate: 73% (95% CI 68–77)
- Symptomatic attack rate among infected: 18% (95% CI 10–27)
- All age groups affected
- Adults more likely to present for medical care
- No severe disease, hospitalizations, or deaths

Note: Rates based on serosurvey on Yap Island, 2007 (population 7,391)



Incubation and viremia

- Incubation period for Zika virus disease is 3–14 days.
- Zika viremia ranges from a few days to 1 week.
- Some infected pregnant women can have evidence of Zika virus in their blood longer than expected.
- Virus remains in semen longer than in blood.



3 - 14 Days




Zika virus clinical disease course and outcomes

- Clinical illness is usually mild.
- Symptoms last several days to a week.
- Severe disease requiring hospitalization is uncommon.
- Fatalities are rare.
- Guillain-Barré syndrome (GBS) reported in patients following suspected Zika virus infection.
 - Relationship to Zika virus infection is not known.



Symptoms

- Many infections asymptomatic.
- Most common symptoms
 - Acute onset of fever
 - Maculopapular rash
 - Joint pain
 - Conjunctivitis
- Other symptoms include muscle pain and headache.



Reported clinical symptoms among confirmed Zika virus disease cases


Symptoms	N (n=31)	%
Macular or papular rash	28	90%
Subjective fever	20	65%
Arthralgia	20	65%
Conjunctivitis	17	55%
Myalgia	15	48%
Headache	14	45%
Retro-orbital pain	12	39%
Edema	6	19%
Vomiting	3	10%

Yap Island, 2007

Duffy M. N Engl J
Med 2009

Clinical Presentation of rash syndrome in Northeast Brazil, May 2015

Rash, swollen joints, and conjunctival hyperemia



Pan American
Health
Organization




World Health
Organization
REGIONAL OFFICE FOR THE Americas

Clinical features: Zika virus compared to dengue and chikungunya

Features	Zika	Dengue	Chikungunya
Fever	++	+++	+++
Rash	+++	+	++
Conjunctivitis	++	-	-
Arthralgia	++	+	+++
Myalgia	+	++	+
Headache	+	++	++
Hemorrhage	-	++	-
Shock	-	+	-

Rabe, Ingrid MBChB, MMed
“Zika Virus- What Clinicians
Need to Know?”
(presentation, Clinician
Outreach and Communication
Activity (COCA) Call, Atlanta,
GA, January 26 2016)

DIAGNOSES AND TESTING FOR ZIKA




Risk of Dengue virus transmission = risk of Zika virus transmission



Differential diagnosis


Based on typical clinical features, the differential diagnosis for Zika virus infection is broad. Considerations include

- Dengue
- Chikungunya
- Leptospirosis
- Malaria
- Rickettsia
- Group A Streptococcus
- Rubella
- Measles
- Parvovirus
- Enterovirus
- Adenovirus
- Other alphaviruses (e.g., Mayaro, Ross River, Barmah Forest, O'nyong-nyong, and Sindbis viruses)



Diagnostic testing for Zika virus

- During first two weeks after the start of illness, Zika virus infection can often be diagnosed by performing real-time reverse transcriptase polymerase chain reaction (rRT-PCR) on serum and urine.
- Serology for IgM and neutralizing antibodies in serum collected up to 12 weeks after illness onset
- Plaque reduction neutralization test (PRNT) for presence of virus-specific neutralizing antibodies in paired serum samples
- Immunohistochemical (IHC) staining for viral antigens or RT-PCR on fixed tissues




Comparison of Test Results for Zika Virus RNA in Urine, Serum, and Saliva Specimens from Persons with Travel-Associated Zika Virus Disease — Florida, 2016

TABLE 1. Results of Zika virus IgM antibody testing of serum specimens and RT-PCR testing of serum and urine specimens for Zika virus RNA, by days after symptom onset for 66 persons with travel-associated Zika virus disease — Florida, 2016

Days after onset	Serum IgM	Serum RT-PCR	Urine
	No. positive/No. tested (%)	No. positive/No. tested (%)	RT-PCR No. positive/No. tested (%)
0	0/1 (0)	0/1 (0)	1/1 (100)
1	2/7 (29)	6/7 (85)	7/7 (100)
2	3/12 (25)	8/12 (67)	11/12 (92)
3	5/10 (50)	4/10 (40)	10/10 (100)
4	3/12 (25)	8/12 (67)	12/12 (100)
5	9/13 (69)	5/13 (38)	11/13 (85)
6	2/2 (100)	0/2 (0)	2/2 (100)
7	4/4 (100)	0/4 (0)	3/4 (75)
9	2/3 (67)	0/3 (0)	3/3 (100)
14	1/1 (100)	0/1 (0)	0/1 (0)
20	1/1 (100)	0/1 (0)	1/1 (100)
Range of days			
0–5	22/55 (40)	31/55 (56)*	52/55 (95)*
6–10	8/9 (89)	0/9 (0)*	8/9 (89)*
11–15	1/1 (100)	0/1 (0)	0/1 (0)
16–20	1/1 (100)	0/1 (0)	1/1 (100)


Serology cross-reactions with other flaviviruses

- Zika virus serology (IgM) can be positive due to antibodies against related *flaviviruses* (e.g., dengue and yellow fever viruses).
- As viremia decreases over time, a negative rRT-PCR collected after symptom onset does not preclude Zika; in this case, serologic testing should be performed.
- Neutralizing antibody testing may discriminate between cross-reacting antibodies in primary *flavivirus* infections.
- Difficult to distinguish infecting virus in people previously infected with or vaccinated against a related *flavivirus*




Laboratories for diagnostic testing

- Testing performed at CDC, select commercial labs, and a few state health departments
- CDC is working to expand laboratory diagnostic testing in states.
- Healthcare providers should work with their state health department to facilitate diagnostic testing and report results.



Recommendations

- CDC recommends Zika virus testing for
 - Symptomatic people who live in or recently traveled to an area with active Zika transmission, and
 - People who have had unprotected sex with someone confirmed to have Zika virus infection or who lives in or traveled to an area with active Zika transmission.
- All pregnant women in the US should be assessed for possible Zika exposure at each prenatal care visit.



Recommendations continued...

- Pregnant women with possible Zika exposure and signs or symptoms consistent with Zika virus disease should be tested based on time of evaluation relative to symptom onset in accordance with [CDC guidance](#).
- Pregnant women with ongoing risk of possible Zika virus exposure and who do not report symptoms of Zika virus disease should be tested in the first and second trimester of pregnancy in accordance with CDC guidance.



REPORTING ZIKA CASES

A solid horizontal bar in a light cyan color, spanning the width of the slide, positioned below the main title.

Reporting cases

- Zika virus disease is a nationally notifiable condition. Report all confirmed cases to your state health department.



Zika pregnancy registry

- CDC established the [US Zika Pregnancy Registry](#) to collect information and learn more about pregnant women in the US with Zika and their infants.
- Data collected will be used to update recommendations for clinical care, plan for services for pregnant women and families affected by Zika, and improve prevention of Zika infection during pregnancy.
- CDC maintains a 24/7 consultation service for health officials and healthcare providers caring for pregnant women. To contact the service, call 770-488-7100 or email ZIKAMCH@cdc.gov
- CDC also established a similar system, the [Zika Active Pregnancy Surveillance System](#), in Puerto Rico.



ZIKA AND PREGNANCY

A solid cyan horizontal bar spans the width of the slide, positioned below the title.

Zika and pregnancy outcomes

- Zika virus can pass from a pregnant woman to her fetus during pregnancy or around the time of birth.
- Zika infection in pregnancy is a cause of microcephaly and other severe brain defects. Other problems include
 - Eye defects, hearing loss, impaired growth, and fetal loss.



Zika and pregnancy outcomes

- Scientists are studying the full range of other potential health problems caused by Zika virus infection during pregnancy.
- No reports of infants getting Zika through breastfeeding.
- No evidence that previous infection will affect future pregnancies.




Who to test for Zika during pregnancy

- All pregnant women should be assessed for Zika at each prenatal care visit. They should be asked if they
 - Traveled to or live in an area with active Zika transmission
 - Had sex without a condom or other barrier method to prevent infection with a partner who lives in or traveled to an area with active Zika transmission



Testing guidance: Pregnant women with possible Zika exposure



Abbreviations: IgM = immunoglobulin M; PRNT = plaque reduction neutralization test; rRT-PCR = real-time reverse transcription–polymerase chain reaction.

* A pregnant woman is considered symptomatic if one or more signs or symptoms (fever, rash, arthralgia, or conjunctivitis) consistent with Zika virus disease is reported whereas a pregnant woman is considered asymptomatic if symptoms are NOT reported.

† Testing includes Zika virus rRT-PCR on serum and urine samples, Zika virus and dengue virus immunoglobulin M (IgM), and plaque reduction neutralization test (PRNT) on serum samples. PRNT results that indicate recent flavivirus infection should be interpreted in the context of the currently circulating flaviviruses. Refer to the laboratory guidance for updated testing recommendations (<http://www.cdc.gov/zika/laboratories/lab-guidance.html>). Because of the overlap of symptoms in areas where other viral illnesses are endemic, evaluate for possible dengue or chikungunya virus infection.

‡ Dengue IgM antibody testing is recommended only for symptomatic pregnant women.

§ If Zika virus rRT-PCR testing is requested from laboratories without IgM antibody testing capacity or a process to forward specimens to another testing laboratory, storing of additional serum samples is recommended for IgM antibody testing in the event of a rRT-PCR negative result.

** Possible exposure to Zika virus includes travel to or residence in an area with active Zika virus transmission (<http://www.cdc.gov/travel/notices>), or sex (vaginal sex (penis-to-vagina sex), anal sex (penis-to-anus sex), oral sex (mouth-to-penis sex or mouth-to-vagina sex), and the sharing of sex toys) without a barrier method to prevent infection (male or female condoms for vaginal or anal sex, male condoms for oral sex (mouth-to-penis), and male condoms cut to create a flat barrier or dental dams for oral sex (mouth-to-vagina) with a partner who traveled to, or lives in an area with active Zika virus transmission).

Clinical management of a pregnant woman with suspected Zika virus infection

Interpretation of Laboratory Results*	Prenatal Management	Postnatal Management
<u>Recent Zika virus infection</u>	<ul style="list-style-type: none"> • Consider serial ultrasounds every 3–4 weeks to assess fetal anatomy and growth[†] • Decisions regarding amniocentesis should be individualized for each clinical circumstance[§] 	<p>LIVE BIRTHS:</p> <ul style="list-style-type: none"> • Cord blood and infant serum should be tested for Zika virus rRT-PCR, Zika IgM, and dengue virus IgM antibodies. If CSF is obtained for other reasons, it can also be tested. • Zika virus rRT-PCR and IHC staining of umbilical cord and placenta is recommended.[†] <p>FETAL LOSSES:</p> <ul style="list-style-type: none"> • Zika virus rRT-PCR and IHC staining of fetal tissues is recommended.[†]
<u>Recent flavivirus infection; specific virus cannot be identified</u>		<ul style="list-style-type: none"> • Consider serial ultrasounds every 3–4 weeks to assess fetal anatomy and growth[†] • Amniocentesis might be considered; decision should be individualized for each clinical circumstance[§]
<u>Presumptive recent Zika virus infection**</u>	<ul style="list-style-type: none"> • Consider serial ultrasounds every 3–4 weeks to assess fetal anatomy and growth[†] • Amniocentesis might be considered; decision should be individualized for each clinical circumstance[§] 	
<u>Presumptive recent flavivirus infection**</u>		<ul style="list-style-type: none"> • Clinical management in accordance with existing guidelines (http://apps.who.int/iris/bitstream/10665/44188/1/9789241547871_eng.pdf).
<u>Recent dengue virus infection</u>	<ul style="list-style-type: none"> • Prenatal ultrasound to evaluate for fetal abnormalities consistent with congenital Zika virus syndrome.[†] • Fetal abnormalities present: repeat Zika virus rRT-PCR and IgM test; base clinical management on corresponding laboratory results. • Fetal abnormalities absent: base obstetric care on the ongoing risk of Zika virus exposure to the pregnant woman. 	
<u>No evidence of Zika virus or dengue virus infection</u>	<ul style="list-style-type: none"> • Prenatal ultrasound to evaluate for fetal abnormalities consistent with congenital Zika virus syndrome.[†] • Fetal abnormalities present: repeat Zika virus rRT-PCR and IgM test; base clinical management on corresponding laboratory results. • Fetal abnormalities absent: base obstetric care on the ongoing risk of Zika virus exposure to the pregnant woman. 	<ul style="list-style-type: none"> • Prenatal ultrasound to evaluate for fetal abnormalities consistent with congenital Zika virus syndrome.[†] • Fetal abnormalities present: repeat Zika virus rRT-PCR and IgM test; base clinical management on corresponding laboratory results. • Fetal abnormalities absent: base obstetric care on the ongoing risk of Zika virus exposure to the pregnant woman.

Zika and pregnancy: Clinical management for obstetricians

- Positive or inconclusive Zika virus testing results
 - Antepartum
 - Consider serial ultrasounds every 3-4 weeks.
 - Referral to maternal-fetal medicine specialist is recommended.
 - Postpartum
 - Histopathologic examination of the placenta and umbilical cord.
 - Testing of frozen placental tissue and cord tissue for Zika virus RNA.
 - Testing of cord serum for Zika and dengue virus IgM and neutralizing antibodies.

**CLINICAL MANAGEMENT OF
INFANTS WITH CONFIRMED
OR POSSIBLE ZIKA INFECTION**




Infants with confirmed or possible Zika infection

Doctors have found problems among fetuses and infants infected with Zika virus before birth, including

- Microcephaly
- Miscarriage
- Stillbirth
- Absent or poorly developed brain structures
- Defects of the eye
- Hearing deficits
- Impaired growth



Testing guidance for infants of women with possible exposure to Zika during pregnancy



*Areas with Zika virus transmission are listed on CDC's website at <http://wwwnc.cdc.gov/travel/notices>.

†Microcephaly defined as occipitofrontal circumference less than the third percentile for gestational age and sex based on standard growth curves, not explained by other etiologies.

‡Laboratory evidence of Zika virus infection includes 1) detectable Zika virus, Zika virus RNA, or Zika virus antigen in any clinical specimen; or 2) positive Zika virus Immunoglobulin M (IgM) with confirmatory neutralizing antibody titers that are ≥ 4 -fold higher than dengue virus neutralizing antibody titers in serum or cerebrospinal fluid. Testing is considered inconclusive if Zika virus neutralizing antibody titers are < 4 -fold higher than dengue virus neutralizing antibody titers.

More information on laboratory testing for Zika virus infection is available at <http://www.cdc.gov/zika/state-labs/index.html>.

Evaluation for all infants with positive or inconclusive Zika virus test results

- Physical examination, measurement of head circumference, and assessment of gestational age
- Evaluation neurologic abnormalities, dysmorphic features, enlarged liver or spleen, and rash/other skin lesions
- Cranial ultrasound
- Ophthalmologic evaluation before hospital discharge or within 1 month after birth
- Evaluation of hearing by evoked otoacoustic emissions testing or auditory brainstem response testing before hospital discharge or within 1 month after birth
- Consultation with appropriate specialist for any abnormal findings

Additional evaluation for infants who have microcephaly or other findings consistent with congenital Zika virus infection

- Consultation with clinical geneticist or dysmorphologist and pediatric neurologist
- Testing for other congenital infections; consider consultation with pediatric infectious disease specialist
- Complete blood count, platelet count, and liver function and enzyme tests
- Genetic or other teratogenic causes should be considered if additional anomalies are identified.

Long term follow up for infants with positive or inconclusive Zika virus test results

- Additional hearing screen at 6 months of age and audiology follow up of abnormal newborn hearing screening
- Continued evaluation of developmental characteristics and milestones, as well as head circumference, through 1st year of life
- Consultation with appropriate medical specialists (e.g., pediatric neurology, developmental and behavioral pediatrics, physical and speech therapy) if any abnormalities are noted and as concerns arise


Case definition of microcephaly

Definite congenital microcephaly for live births

- Head circumference (HC) at birth is less than the 3rd percentile for gestational age and sex.
- If HC at birth is not available, HC less than the 3rd percentile for age and sex within the first 6 weeks of life.

Definite congenital microcephaly for still births and early termination

- HC at delivery is less than the 3rd percentile for gestational age and sex.



Definitions for *possible* congenital microcephaly

Possible congenital microcephaly for live births


- If earlier HC is not available, HC less than 3rd percentile for age and sex beyond 6 weeks of life.

Possible microcephaly for all birth outcomes

- Microcephaly diagnosed or suspected on prenatal ultrasound in the absence of available HC measurements.



Measuring head circumference for microcephaly



- Use a measuring tape that cannot be stretched
- Securely wrap the tape around the widest possible circumference of the head
 - Broadest part of the forehead above eyebrow
 - Above the ears
 - Most prominent part of the back of the head
- Take the measurement three times and select the largest measurement to the nearest 0.1 cm
- Optimal measurement at 24-36 hours after birth when molding of the head has subsided

SEXUAL TRANSMISSION

A solid cyan horizontal bar spans the width of the slide, positioned below the title.

Centers for Disease Control and Prevention

MMWR

Morbidity and Mortality Weekly Report

Weekly / Vol. 65 / No. 28

July 22, 2016

Suspected Female-to-Male Sexual Transmission of Zika Virus — New York City, 2016



Pan American
Health
Organization



REGIONAL OFFICE FOR THE

World Health
Organization
Americas

About sexual transmission

- Zika virus can be passed through sex, even if the infected person does not have symptoms at the time.
- Sex includes vaginal, anal, and oral sex, and the sharing of sex toys.
- Condoms and other barriers* can reduce the chance of getting Zika from sex.
- Not having sex can eliminate the risk of getting Zika from sex.

* Barriers include male and female condoms and dental dams.



What we do not know about sexual transmission

- We do not know how often people with Zika who never develop symptoms pass Zika through sex.
- We do not know if sexual transmission of Zika virus poses a different risk of birth defects than mosquito-borne transmission.



Preventing or reducing the chance of sexual transmission for couples who are pregnant

- Not having sex can eliminate the risk of getting Zika from sex.
- Condoms can reduce the chance of getting Zika from sex.
- Pregnant couples with a partner who lives in or recently traveled to an area with Zika should use condoms (or other barriers to prevent infection) correctly every time they have sex or not have sex during pregnancy.




Non-pregnant couples with a partner who traveled to an area with Zika

- For non-pregnant couples with a partner who has **recently traveled** to an area with Zika
 - At least 8 weeks after a Zika diagnosis or start of symptoms if the traveling partner is female or if the traveling partner (male or female) has no symptoms.
 - At least 6 months after a Zika diagnosis or start of symptoms if the traveling partner is male. This long extended period is because Zika stays in semen longer than in other body fluids.



Non-pregnant couples with a partner who lives in an area with Zika

- Couples living in an area with Zika can use condoms or not have sex as long as there is Zika in the area. If either partner develops symptoms of Zika or has concerns, they should talk to a healthcare provider.



Update: Interim Guidance for Prevention of Sexual Transmission of Zika Virus — United States, July 2016

BOX. Recommendations for prevention of sexual transmission of Zika virus for couples in which one or both partners have traveled to or reside in an area with active Zika virus transmission

Couples in which a woman is pregnant

- Couples in which a woman is pregnant should use barrier methods against infection consistently and correctly or abstain from sex for the duration of the pregnancy.

Couples who are not pregnant and are not planning to become pregnant*

- Couples in which a partner had confirmed Zika virus infection or clinical illness consistent with Zika virus disease should consider using barrier methods against infection consistently and correctly or abstain from sex as follows:
 - Men with Zika virus infection for at least 6 months after onset of illness;
 - Women with Zika virus infection for at least 8 weeks after onset of illness.
- Couples in areas without active Zika transmission in which one partner traveled to or resides in an area with active Zika virus transmission but did not develop symptoms of Zika virus disease should consider using barrier methods against infection or abstaining from sex for at least 8 weeks after that partner departed the Zika-affected area.
- Couples who reside in an area with active Zika virus transmission might consider using barrier methods against infection or abstaining from sex while active transmission persists.

PRECONCEPTION GUIDANCE

A solid horizontal bar in a light cyan color, spanning the width of the slide, positioned below the main title.

Couples interested in conceiving who DO NOT reside in an area with active Zika virus transmission

- For Women with possible exposure to Zika virus
 - Discuss signs and symptoms and potential adverse outcomes associated with Zika
 - If Zika virus disease diagnosed or symptoms develop, wait at least 8 weeks after symptom onset to attempt conception.
 - If NO symptoms develop, wait at least 8 weeks after last date of exposure before attempting conception.
 - During that time, use condoms every time during sex or do not have sex to protect partner.



Couples interested in conceiving who DO NOT reside in an area with active Zika virus transmission

- For Men with possible exposure to Zika virus
 - If Zika virus disease diagnosed or symptoms develop, wait at least 6 months after symptom onset to attempt conception.
 - If NO symptoms develop, wait at least 8 weeks after exposure to attempt conception.
 - During that time, use condoms every time during sex or do not have sex to protect partner.
 - Discuss contraception and use of condoms.



Couples interested in conceiving who reside in an area with active Zika virus transmission

- Women and men interested in conceiving should talk with their HCPs
- Factors that may aid in decision-making
 - Reproductive life plan
 - Environmental risk of exposure
 - Personal measures to prevent mosquito bites
 - Personal measures to prevent sexual transmission
 - Education about Zika virus infection in pregnancy
 - Risks and benefits of pregnancy at this time




WHAT TO TELL PATIENTS ABOUT ZIKA

A solid cyan horizontal bar spans the width of the slide, positioned below the title.

Pregnant women

- Should not travel to areas with Zika.
- If they must travel to areas with Zika, tell pregnant patients to protect themselves from mosquito bites and take steps to prevent sexual transmission during and after travel.




Treating patients who test positive


- There are no vaccine or medicine Zika.
- Treat the symptoms of Zika
 - Rest
 - Drink fluids to prevent dehydration
 - Take acetaminophen (Tylenol®) to reduce fever and pain
 - Do not take aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs) until dengue can be ruled out to reduce the risk of bleeding.



Patients who test positive

- Protect from mosquito bites during the first week of illness, when Zika virus can be found in blood.
- The virus can be passed from an infected person to a mosquito through bites.
- An infected mosquito can spread the virus to other people.





Projected Zika Virus Importation and Subsequent Ongoing Transmission after Travel to the 2016 Olympic and Paralympic Games — Country-Specific Assessment, July 2016

BOX. CDC prevention recommendations for athletes and visitors to Rio de Janeiro, and other areas where Zika virus is circulating

- Pregnant women should not travel to any area where Zika virus transmission is ongoing.
- Travelers should take protective measures, including use of insect repellent, to prevent mosquito bites both during travel and for 3 weeks after returning to their home country. Such measures include wearing long-sleeved shirts and long pants; staying in places with air conditioning and window and door screens to keep mosquitoes outside; sleeping under a mosquito bed net, and using insect repellents with active ingredients (e.g., DEET).
- Travelers should prevent possible sexual transmission while at the 2016 Olympic and Paralympic Games and after returning home by correctly using condoms every time they have sex or by abstaining from sex. Males should use condoms for at least 8 weeks after travel or, if symptomatic for Zika virus infection, for 6 months from the start of symptoms.
- After returning from a country with Zika virus transmission, men with pregnant partners should use condoms or not have sex for the duration of the pregnancy.
- Couples who want to try to get pregnant after attending the Olympic and Paralympic Games should wait at least 8 weeks, and 6 months if the male partner has symptomatic Zika virus infection.


TABLE 2. Participating countries currently not reporting Zika outbreaks (n = 19) that met risk criteria for Zika virus importation and subsequent ongoing transmission attributed to travel to the Olympic and Paralympic Games, ranked by aviation travel volume* from Rio de Janeiro, Brazil — August 2016

Country	Total aviation travel from Rio de Janeiro in August 2015 (no. passenger-journeys)	No. of athletes for 2016 Games [†]	Estimated size of Olympic delegation for countries without aviation travel from Rio de Janeiro in August 2015 (no. persons)	Aviation travel from all Zika-affected countries in 2015 [§] No. passengers (% travel attributable to Rio de Janeiro [¶])
Angola	2,841	21	NA	87,549 (3.25)
China	1,201	379	NA	308,238 (0.39)
Hong Kong**	229	32	NA	108,215 (0.21)
Sao Tome and Principe	104	2	NA	4,732 (2.20)
Oman	70	3	NA	2,300 (3.04)
Saudi Arabia	46	8	NA	7,688 (0.60)
Congo	22	6	NA	2,023 (1.09)
Myanmar	15	1	NA	733 (2.05)
Antigua and Barbuda	10	3	NA	61,312 (0.02)
Cayman Islands	10	3	NA	66,859 (0.01)
Ghana	10	6	NA	2,498 (0.40)
Rwanda	1	5	—	281 (0.36)
Eritrea	0	9	27	27 (100)
Yemen	0	4 ^{††}	12	13 (92)
Djibouti	0	4	12	62 (19.35)
The Gambia	0	3	9	1,198 (0.75)
Chad	0	2 ^{††}	6	19(31.6)
Mauritania	0	2 ^{††}	6	201 (2.99)
Sudan	0	1	3	153 (1.96)

WHAT TO TELL PATIENTS ABOUT MOSQUITO BITE PROTECTION

Mosquito bite protection

- Wear long-sleeved shirts and long pants.
- Stay and sleep in places with air conditioning and window and door screens to keep mosquitoes outside.
- Take steps to [control mosquitoes inside and outside your home](http://www.cdc.gov/zika/prevention/controlling-mosquitoes-at-home.html) (<http://www.cdc.gov/zika/prevention/controlling-mosquitoes-at-home.html>).
- Sleep under a mosquito bed net if you are overseas or outside and are not able to protect yourself from mosquito bites.



Mosquito bite protection

- Use Environmental Protection Agency (EPA)-registered insect repellents with one of the following active ingredients: DEET, picaridin, IR3535, oil of lemon eucalyptus, or para-menthane-diol.
- Always follow the product label instructions.
- Do not spray repellent on the skin under clothing.
- If you are also using sunscreen, apply sunscreen before applying insect repellent.



Mosquito bite protection

- Do not use insect repellent on babies younger than 2 months old.
- Do not use products containing oil of lemon eucalyptus or para-menthane-diol on children younger than 3 years old.
- Dress children in clothing that covers arms and legs.
- Cover crib, stroller, and baby carrier with mosquito netting.
- Do not apply insect repellent onto a child's hands, eyes, mouth, and cut or irritated skin.
 - Adults: Spray insect repellent onto your hands and then apply to a child's face.



The Medical Letter[®]

on Drugs and Therapeutics

Insect Repellents

Table 1. Some Insect Repellents

Repellent	Brand Name	Formulation	Duration of Protection ¹		Cost (Size) ²
			Mosquitoes	Ticks	
DEET	<i>Cutter Skinsations</i>	7% pump spray	1-3 hrs ³	6 hrs ³	\$6.40 (7.5 oz)
	<i>Repel Scented Family</i>	15% aerosol spray	5-8 hrs ³	8.5 hrs ³	6.86 (6.5 oz)
	<i>Off Deep Woods VIII</i>	25% aerosol spray	8 hrs ³	5 hrs ³	7.38 (6 oz)
	<i>Sawyer Ultra 30 Liposome Controlled Release</i>	30% lotion	11 hrs	N.A.	8.49 (4 oz)
	<i>Ultrathon⁴</i>	34% lotion	12 hrs	N.A.	6.99 (2 oz)
Picaridin	<i>Cutter Advanced</i>	5.75% wipes	8 hrs	5 hrs	5.99 (18 wipes)
	<i>Avon Skin So Soft Bug Guard Plus Picaridin</i>	10% aerosol spray	8 hrs	12 hrs	6.99 (4 oz)
	<i>Natrapel 8 hour</i>	20% pump spray	8 hrs	8 hrs	5.59 (3.4 oz)
IR3535	<i>Avon Skin So Soft Bug Guard Plus IR3535⁵</i>	7.5% lotion	2 hrs	2 hrs	10.89 (4 oz)
	<i>Coleman Skin Smart</i>	20% pump spray	8 hrs	8 hrs	4.99 (5 oz)
	Oil of lemon eucalyptus	<i>Coleman Botanicals</i>	30% pump spray	6 hrs	N.A.
<i>Repel Lemon Eucalyptus</i>		30% pump spray	7-8 hrs ³	7 hrs ³	4.99 (4 oz)
Permethrin	<i>Sawyer Premium Permethrin Clothing</i>	0.5% pump spray	—	—	14.99 (24 oz)
	<i>Repel Permethrin Clothing and Gear</i>	0.5% aerosol spray	—	—	7.42 (6.5 oz)

“There are risks and costs to action. But they are far less than the long range risks of comfortable inaction”.

John F. Kennedy